

FUZZY ABEL GRASSMANN'S GROUPOIDS

MADAD KHAN AND M. NOUMAN ASLAM KHAN

ABSTRACT. In the present paper we have studied the concept of fuzzification in AG-groupoids. The equivalent statement for an AG-groupoid to be a commutative semigroup is proved. Fuzzy points have been defined in an AG-groupoid and has been shown the representation of smallest fuzzy left ideal generated by a fuzzy point. The set of all fuzzy left ideals, which are idempotents, forms a commutative monoid. The relation of fuzzy left(right) ideals, fuzzy interior ideals and fuzzy bi-ideals in AG-groupoid has been studied. Necessary and sufficient condition of fully fuzzy prime AG-groupoid has been shown. Further, It has been shown that the set of fuzzy quasi-prime ideals of AG-groupoid with left identity forms a semilattice structure. Moreover, equivalent statements for fuzzy semiprime left ideal in an AG-groupoid have been proved.

1. Introduction

A fuzzy subset f of a set S is a function from S to a closed interval $[0, 1]$. The concept of a fuzzy subset of a set was first considered by L.A. Zadeh [10] in 1965. A. Rosenfelt was the first who studied fuzzy sets in the structure of groups [11]. Fuzzy semigroups were first considered by N. Kuroki [12] in which he studied the bi-ideals in semigroups. The fuzzy theory provides the underlying structure for the generalization of many fields including logic, differential equations and groups. The fuzzy theory on algebraic structures have been widely explored. An Abel Grassmann's groupoid, abbreviated as AG-groupoid, is a groupoid S whose elements satisfy the left invertive law: $(ab)c = (cb)a$ for all a, b and c in S . An AG-groupoid is the midway structure between a commutative semigroup and a groupoid. It is a useful non-associative structure with wide applications in theory of flocks. This property of an AG-groupoid inspired us to use the fuzzy theory in AG-groupoids. In an AG-groupoid the medial law, $(ab)(cd) = (ac)(bd)$ for all a, b, c and d in S holds [6]. If there exists an element e in an AG-groupoid S such that $ex = x$ for all x in S then S is called an AG-groupoid with left identity e . It is to be noted that if an AG-groupoid S has the right identity then S is a commutative monoid. If an AG-groupoid S contains left identity then $(ab)(cd) = (dc)(ba)$ holds for all a, b, c and d in S . Also $a(bc) = b(ac)$ holds for all a, b and c in an AG-groupoid with left identity. In an AG-groupoid S , an element $a \in S$ is called idempotent if $a^2 = a$.

Let $F(S)$ denote the collection of all fuzzy subsets of S . For subsets A, B of S , $AB = \{ab \in S : a \in A, b \in B\}$. A non-empty subset A of S is called left(right) ideal of S if $SA \subseteq A$ ($AS \subseteq A$). Further A is called two-sided ideal if it is both left and right ideal of S . A non-empty subset A of S is called interior ideal of S if $(SA)S \subseteq A$. A non-empty AG-subgroupoid A of S is called bi-ideal of S if

2000 *Mathematics Subject Classification.* 20M10, 20N990.

Key words and phrases. Fuzzy ideal, AG-groupoid, Fuzzy left ideal generated by fuzzy points.

$(AS)A \subseteq A$. A non-empty subset A of S is called idempotent if $AA = A$. An ideal P of S is called prime if $AB \subseteq P$ implies that either $A \subseteq P$ or $B \subseteq P$, for all ideals A and B of S . An AG-groupoid S is called fully prime if every ideal is prime. The left ideal P of S is called quasi-prime if $AB \subseteq P$ implies that either $A \subseteq P$ or $B \subseteq P$, for all left ideals A and B of S . An AG-groupoid S is called fully quasi-prime if every left ideal is quasi-prime.

2. FUZZY IDEALS IN AG-GROUPOIDS

Let f and g be two fuzzy subsets of an AG-groupoid S . The product $f \circ g$ is defined by

$$(f \circ g)(x) = \begin{cases} \bigvee_{x=yz} \{f(y) \wedge g(z)\}, & \text{if } \exists y \text{ and } z \in S, \text{ such that } x = yz, \\ 0 & \text{otherwise.} \end{cases}$$

Proposition 1. *Let S be an AG-groupoid, then the set $(F(S), \circ)$ is an AG-groupoid.*

Proof. Clearly $F(S)$ is closed. Let f, g and h be in $F(S)$. Then for any x in S we have, $((f \circ g) \circ h)(x) = \bigvee_{x=yz} \{(f \circ g)(y) \wedge h(z)\} = \bigvee_{x=yz} \left\{ \bigvee_{y=pq} \{f(p) \wedge g(q)\} \wedge h(z) \right\} = \bigvee_{x=(pq)z} \{f(p) \wedge g(q) \wedge h(z)\} = \bigvee_{x=(zq)p} \{h(z) \wedge g(q) \wedge f(p)\}$. Then, further we have $\bigvee_{x=wp} \left\{ \bigvee_{w=zq} \{h(z) \wedge g(q) \wedge f(p)\} \right\} = \bigvee_{x=wp} \{(h \circ g)(w) \wedge f(p)\} = ((h \circ g) \circ f)(x)$. Hence $(F(S), \circ)$ is an AG-groupoid. $\square$

Corollary 1. *Let S be an AG-groupoid, then the medial law holds in $F(S)$.*

Proof. Let f, g, h , and k be arbitrary elements of $F(S)$. By successive use of left invertive law, $(f \circ g) \circ (h \circ k) = ((h \circ k) \circ g) \circ f = ((g \circ k) \circ h) \circ f = (f \circ h) \circ (g \circ k)$. $\square$

Theorem 1. *Let S be an AG-groupoid with left identity, then the following properties hold in $F(S)$;*

- (i) $f \circ (g \circ h) = g \circ (f \circ h)$ for all f, g and h in $F(S)$,
- (ii) $(f \circ g) \circ (h \circ k) = (k \circ h) \circ (g \circ f)$ for all f, g, h and k in $F(S)$.

Proof. (i) Let x be an arbitrary element of S . If x is not expressible as a product of two elements in S , then $(f \circ (g \circ h))(x) = 0 = (g \circ (f \circ h))(x)$. Let there exists y and z in S such that $x = yz$, then $(f \circ (g \circ h))(x) = \bigvee_{x=yz} \{f(y) \wedge (g \circ h)(z)\} = \bigvee_{x=yz} \left\{ f(y) \wedge \bigvee_{z=pq} \{g(p) \wedge h(q)\} \right\} = \bigvee_{x=y(pq)} \{f(y) \wedge g(p) \wedge h(q)\}$, which implies that $\bigvee_{x=p(yq)} \{g(p) \wedge f(y) \wedge h(q)\} = \bigvee_{x=pw} \left\{ g(p) \wedge \bigvee_{w=yq} \{f(y) \wedge h(q)\} \right\}$ and so we have $\bigvee_{x=pw} \{g(p) \wedge (f \circ h)(w)\} = (g \circ (f \circ h))(x)$. Thus, $(f \circ (g \circ h))(x) = (g \circ (f \circ h))(x)$. If z is not expressible as a product of two elements in S , then $(f \circ (g \circ h))(x) = 0 = (g \circ (f \circ h))(x)$. Hence, $(f \circ (g \circ h))(x) = (g \circ (f \circ h))(x)$ for all x in S .

(ii) If any element x of S is not expressible as product of two elements in S at any stage, then $((f \circ g) \circ (h \circ k))(x) = 0 = ((k \circ h) \circ (g \circ f))(x)$. Let there exists y, z in S such that $x = yz$, then $((f \circ g) \circ (h \circ k))(x) = \bigvee_{x=yz} \{(f \circ g)(y) \wedge (h \circ k)(z)\} =$
 $\bigvee_{x=yz} \left\{ \bigvee_{y=pq} \{f(p) \wedge g(q)\} \wedge \bigvee_{z=uv} \{h(u) \wedge k(v)\} \right\}$ and hence the above equality becomes $\bigvee_{x=(pq)(uv)} \{f(p) \wedge g(q) \wedge h(u) \wedge k(v)\} = \bigvee_{x=(vu)(qp)} \{k(v) \wedge h(u) \wedge g(q) \wedge f(p)\} =$
 $\bigvee_{x=mn} \left\{ \bigvee_{m=vu} \{k(v) \wedge h(u)\} \wedge \bigvee_{n=qp} \{g(q) \wedge f(p)\} \right\} = \bigvee_{x=mn} \{(k \circ h)(m) \wedge (g \circ f)(n)\} =$
 $((k \circ h) \circ (g \circ f))(x).$ $\square$

Proposition 2. *An AG-groupoid S with $F(S) = (F(S))^2$ is commutative semigroup if and only if $(f \circ g) \circ h = f \circ (h \circ g)$ holds for all fuzzy subsets f, g and h of S .*

Proof. Let an AG-groupoid S is commutative semigroup. For any fuzzy subsets f, g and h of S by use of proposition 1 and commutative law, $(f \circ g) \circ h = (h \circ g) \circ f = h \circ (f \circ g)$. Conversely let $(f \circ g) \circ h = f \circ (h \circ g)$ holds for all fuzzy subsets f, g and h of S . We have to show that an AG-groupoid S is commutative semigroup. Let f and g be any arbitrary fuzzy subsets of S . Since $F(S) = (F(S))^2$ so $f = h \circ k$ where h and k are any fuzzy subsets of S . Now $f \circ g = (h \circ k) \circ g = (g \circ k) \circ h = g \circ (h \circ k) = g \circ f$, which shows that commutative law holds in S . By successive use of proposition 1 and commutative law $(f \circ g) \circ k = (k \circ g) \circ f = f \circ (k \circ g) = f \circ (g \circ k)$. $\square$

A fuzzy subset of S is called fuzzy AG-subgroupoid of S if $f(ab) \geq f(a) \wedge f(b)$ for all a and b in S , and is called fuzzy left(right) ideal of S if $f(ab) \geq f(b)$ ($f(ab) \geq f(a)$) for all a and b in S . A fuzzy subset f of S is called a fuzzy two sided ideal(or a fuzzy ideal) of S if it is both fuzzy left and fuzzy right ideal of S . A fuzzy subset f of S is called fuzzy idempotent if $f \circ f = f$. For a subset A of S the characteristic function, C_A is defined by $C_A = \begin{cases} 1 & \text{if } x \in A, \\ 0 & \text{if } x \notin A. \end{cases}$

Note that an AG-groupoid S can be considered a fuzzy subset of itself and we write $S = \mathbf{C}_S$, i.e., $S(x) = 1$, for all x in S . Let a be an arbitrary element of S , then for λ in $(0, 1]$ and for x in S we define fuzzy point a_λ of S as; $a_\lambda(x) = \begin{cases} \lambda & \text{if } x = a, \\ 0 & \text{otherwise.} \end{cases}$. A very simple calculations shows that for a fuzzy subset f of S we have $f = \bigcup_{a_\lambda \in f} a_\lambda$.

It is easy to see from [1] that a non-empty subset A of an AG-groupoid S is AG-subgroupoid if and only if C_A is a fuzzy AG-subgroupoid of S , and A is left(right, two-sided) ideal of S if and only if C_A is a fuzzy left(right, two-sided) ideal of S . For non-empty sets A and B of an AG-groupoid S , $C_A \cap C_B = C_{A \cap B}$ and $C_A \circ C_B = C_{AB}$. It is easy to see that for every fuzzy subset f of an AG-groupoid S , we have $f \subseteq S$. The following lemmas have the same proof as in [1].

Lemma 1. *Let f be a fuzzy subset of an AG-groupoid S , then the following properties hold.*

- (i) f is a fuzzy AG-subgroupoid of S if and only if $f \circ f \subseteq f$.
- (ii) f is a fuzzy left ideal of S if and only if $S \circ f \subseteq f$.
- (iii) f is a fuzzy right ideal of S if and only if $f \circ S \subseteq f$.

(iv) f is a fuzzy ideal of S if and only if $S \circ f \subseteq f$ and $f \circ S \subseteq f$.

Lemma 2. *Let S be an AG-groupoid. Then the following properties hold.*

(i) *Let f and g be two fuzzy AG-subgroupoids of S . Then $f \cap g$ is also a fuzzy AG-subgroupoid of S .*

(ii) *Let f and g be two fuzzy left(right, two-sided) ideal of S . Then $f \cap g$ is also a fuzzy left(right, two-sided) ideal of S .*

Lemma 3. *In an AG-groupoid with left identity $S \circ S = S$.*

Proof. Every x in S can be written as $x = ex$, where e is the left identity in S . So $S \circ S(x) = \bigvee_{x=yz} \{S(y) \wedge S(z)\} \geq \{S(e) \wedge S(x)\} = 1$. Hence $(S \circ S)(x) = 1 = S(x)$ for all x in S . $\square$

Lemma 4. *In an AG-groupoid S with left identity, for every fuzzy left ideal f of S , we have $(S \circ f) = f$.*

Proof. It is sufficient to show that $f \subseteq S \circ f$. Now for any x in S , $(S \circ f)(x) = \bigvee_{x=yz} \{S(y) \wedge f(z)\}$. Since $x = ex$, for all x in S , as e is left identity in S , so $\bigvee_{x=yz} \{S(y) \wedge f(z)\} \geq S(e) \wedge f(x) = f(x)$. $\square$

Proposition 3. *Let S be an AG-groupoid with left identity and f and k are fuzzy left ideals in S then for any fuzzy subsets g and h of S , $f \circ g = h \circ k$ implies $g \circ f = k \circ h$.*

Proof. Since f and h are fuzzy left ideals in S so by above lemma $S \circ f = f$ and $S \circ h = h$. Now $g \circ f = (S \circ g) \circ f = (f \circ g) \circ S = (h \circ k) \circ S = (S \circ k) \circ h = k \circ h$. $\square$

The following corollary is direct consequence of the successive use of left invertive law in fuzzy AG-groupoid shown in proposition 1.

Corollary 2. *In an AG-groupoid S for any fuzzy subsets f, g and h the following conditions are equivalent:*

- (i) $(f \circ g) \circ h = g \circ (f \circ h)$,
- (ii) $(f \circ g) \circ h = g \circ (h \circ f)$.

Theorem 2. *If S is an AG-groupoid then $Q = \{f \mid f \in S, f \circ h = f \text{ where } h = h \circ h\}$ is a commutative monoid in S .*

Proof. The subset Q is non-empty since $h \circ h = h$ which implies that h is in Q . Let f and g be fuzzy subsets of S in Q then $f \circ h = f$ and $g \circ h = g$. Consider $f \circ g = (f \circ h) \circ (g \circ h) = (f \circ g) \circ (h \circ h) = (f \circ g) \circ h$ which implies that Q is closed. Now for any fuzzy subsets f and g in Q , $f \circ g = (f \circ h) \circ g = (g \circ h) \circ f = g \circ f$ which implies that commutative law holds in Q and associative law holds in Q due to commutativity. Since for any fuzzy subset f in Q we have $f \circ h = f$, where h is fixed, implies that h is a right identity in S and hence an identity. $\square$

Let S be an AG-groupoid and a_λ be a fuzzy point in S . The smallest fuzzy left ideal of S containing a_λ is called a fuzzy left ideal of S generated by a_λ . Let we denote the smallest fuzzy left ideal of S generated by a_λ by $\langle a_\lambda \rangle_L$.

Theorem 3. *Let S be an AG-groupoid with left identity e and a_λ be a fuzzy point in S , then $\langle a_\lambda \rangle_L = f$, where $f(x) = \begin{cases} \lambda & \text{if there exists } b \in S \text{ such that } x = ba, \\ 0 & \text{otherwise.} \end{cases}$*

Proof. Let x and y be an arbitrary elements of S . Let $f(y) = 0$ then $f(xy) \geq 0 = f(y)$ and if $f(y) \neq 0$ then there exists an element b in S such that $y = ba$. Now, $f(xy) = f(x(ba)) = f((ex)(ba)) = f((ab)(xe)) = f(((xe)b)a) = \lambda \geq f(y)$, which implies that f is a fuzzy left ideal in S . Since $a = ea$, so $f(a) = f(ea) = \lambda$ and hence a_λ is in f . Let g be any other left ideal of S containing a_λ , then $g(a) \geq \lambda$. Let $f(x) = 0$ for all x in S then $g(x) \geq 0 = f(x)$. On the other hand if $f(x) = \lambda$ then there exists b in S such that $x = ba$ and $g(x) = g(ba) \geq g(a) \geq \lambda = f(x)$, which implies that $f \subseteq g$. Hence f is the smallest left ideal generated by a in S . $\square$

Proposition 4. *Let S be an AG-groupoid, then every fuzzy left ideal which is idempotent is a fuzzy ideal.*

Proof. Let f be a fuzzy left ideal in S , which is idempotent. Consider, $f \circ S = (f \circ f) \circ S = (S \circ f) \circ f \subseteq f \circ f = f$. $\square$

Remark 1. *If S is an AG-groupoid with left identity, then in above proposition fuzzy left ideal and fuzzy right ideal coincide.*

Theorem 4. *Let f is fuzzy idempotent in AG-groupoid S with left identity, then followings are true.*

- (i) $S \circ f$ is an idempotent.
- (ii) Every fuzzy left ideal g in S commutes with f .

Proof. (i) can easily be shown by use of corollary 1 and lemma 3.

For (ii) consider, $f \circ g = (f \circ f) \circ g = (g \circ f) \circ f \subseteq (g \circ S) \circ f \subseteq g \circ f$. Also, $g \circ f = g \circ (f \circ f) = f \circ (g \circ f) \subseteq f \circ (g \circ S) \subseteq f \circ g$. $\square$

Theorem 5. *Let S be an AG-groupoid with left identity, then the collection of all fuzzy left ideals of S , which are idempotent forms a commutative monoid.*

Proof. Let $\hat{H}$ denote the collection of all fuzzy left ideals which are idempotent in S . Here, $\hat{H}$ is non-empty, since by lemma 3, $S \circ S = S$ implies S is in $\hat{H}$. Consider f, g in $\hat{H}$, then $(f \circ g) \circ (f \circ g) = (f \circ f) \circ (g \circ g) = f \circ g$, also by corollary 1, $S \circ (f \circ g) = (S \circ S) \circ (f \circ g) = (S \circ f) \circ (S \circ g) \subseteq f \circ g$. Also for every f, g in $\hat{H}$ by use of theorem 1 and corollary 1, $f \circ g = (f \circ g) \circ (f \circ g) = (g \circ f) \circ (g \circ f) = (g \circ g) \circ (f \circ f) = g \circ f$, that is commutative law holds in $\hat{H}$. Now, for any f, g , and h in $\hat{H}$ we have $(f \circ g) \circ h = (h \circ g) \circ f = f \circ (h \circ g) = f \circ (g \circ h)$. Since every f in $\hat{H}$ is left ideal, so lemma 4 implies that $S \circ f = f$. Commutativity implies that $S \circ f = f \circ S = f$, which implies that S is identity in $\hat{H}$ and every f is an ideal in $\hat{H}$. $\square$

Lemma 5. *Let S be an AG-groupoid with left identity e . Then every fuzzy right ideal is fuzzy ideal.*

Proof. Let f be a fuzzy right ideal in S , so $f \circ S \subseteq f$. By use of lemma 3 and proposition 1, $S \circ f = (S \circ S) \circ f = (f \circ S) \circ S \subseteq f \circ S \subseteq f$. So f is a fuzzy left ideal, and hence a fuzzy ideal in S . $\square$

Remark 2. *If f is a fuzzy right ideal of an AG-groupoid S with left identity then $f \cup (S \circ f)$ and $f \cup (f \circ f)$ are fuzzy two-sided ideals of S .*

Lemma 6. *If f is a fuzzy left ideal of an AG-groupoid S with left identity then $f \cup (f \circ S)$ and $f \cup (f \circ f)$ are fuzzy two-sided ideals of S .*

Proof. Consider, $(f \cup (f \circ S)) \circ S = (f \circ S) \cup ((f \circ S) \circ S) = (f \circ S) \cup ((S \circ S) \circ f) = (f \circ S) \cup (S \circ f) = (f \circ S) \cup f = f \cup (f \circ S)$. Hence $f \cup (f \circ S)$ is fuzzy right ideal of S , and by lemma 5, $f \cup (f \circ S)$ is fuzzy two-sided ideal of S . Now $(f \cup (f \circ f)) \circ S = (f \circ S) \cup (f \circ f) \circ S = (f \circ S) \cup ((S \circ f) \circ f) \subseteq (f \circ S) \cup (f \circ f) = (f \circ f) \cup (S \circ f) \subseteq (f \circ f) \cup f = f \cup (f \circ f)$, implies that $f \cup (f \circ f)$ is fuzzy right ideal of S . By lemma 5, $f \cup (f \circ f)$ is fuzzy left ideal of S . $\square$

A fuzzy subset f of an AG-groupoid S is called a fuzzy bi-ideal of S if $f((xy)z) \geq f(x) \wedge f(z)$ for all x, y and z of S . It is easy to see from [1] that for a non-empty subset A of an AG-groupoid S is a bi-ideal of S if and only if C_A is a fuzzy bi-ideal of S . The following lemma has proof as in [1].

Lemma 7. *Let f be a fuzzy AG-subgroupoid of an AG-groupoid S . Then f is a fuzzy bi-ideal of S if and only if $(f \circ S) \circ f \subseteq f$.*

Lemma 8. *Let f and g be fuzzy right ideals of an AG-groupoid S with left identity. Then $f \circ g$ and $g \circ f$ are fuzzy bi-ideals of S .*

Proof. By use of corollary 1 $(f \circ g) \circ (f \circ g) = (f \circ f) \circ (g \circ g) \subseteq f \circ g$. Hence $f \circ g$ is fuzzy AG-subgroupoid of S . Now, by proposition 1, lemma 3 and corollary 1 $((f \circ g) \circ S) \circ (f \circ g) = ((f \circ g) \circ (S \circ S)) \circ (f \circ g) = ((f \circ S) \circ (g \circ S)) \circ (f \circ g) \subseteq (f \circ g) \circ (f \circ g) \subseteq f \circ g$. Similarly, $g \circ f$ is a bi-ideal. $\square$

Lemma 9. *Let f and g be fuzzy bi-ideals of an AG-groupoid S . Then $f \cap g$ is a fuzzy bi-ideal of S .*

Proof. It is same as in [1]. $\square$

A fuzzy subset f of an AG-groupoid S is called a fuzzy interior ideal of S if $f((xa)y) \geq f(a)$ for all x, a and y of S . It can easily seen that, if A be a non-empty subset of an AG-groupoid S , then A is a interior ideal of S if and only if C_A is a fuzzy interior ideal of S . The following lemma has the proof as in [1].

Lemma 10. *Let f be a fuzzy AG-subgroupoid of an AG-groupoid S . Then f is a fuzzy interior ideal of S if and only if $(S \circ f) \circ S \subseteq f$.*

Proposition 5. *Let S be an AG-groupoid. Then for any fuzzy left ideal, which is idempotent, in S , the following properties hold.*

- (i) f is a fuzzy bi-ideal.
- (ii) f is a fuzzy interior ideal.

Proof. (i) Since a fuzzy subset f of S is fuzzy left ideal so $f \circ f \subseteq f$. By use of corollary 1 $(f \circ S) \circ f = (f \circ S) \circ (f \circ f) = (f \circ f) \circ (S \circ f) \subseteq f \circ f = f$.

(ii) Consider, $(S \circ f) \circ S \subseteq f \circ S = (f \circ f) \circ S = (S \circ f) \circ f \subseteq f \circ f = f$, implies that f is an interior ideal of S . $\square$

Lemma 11. *Every fuzzy subset f of an AG-groupoid S with left identity is right ideal if and only if f is an interior ideal.*

Proof. Let every fuzzy subset f of S is right ideal. For x, a and y of S , consider $f((xa)y) \geq f(xa) = f((ex)a) = f((ax)e) \geq f(ax) \geq f(a)$, which implies that f is an interior ideal. Conversely, for any x and y in S we have, $f(xy) = f((ex)y) \geq f(x)$. $\square$

Lemma 12. *Let f be a fuzzy left ideal in an AG-groupoid S with left identity, then f being interior ideal is bi-ideal of S .*

Proof. Since f is fuzzy left ideal in S , so $f(xy) \geq f(y)$ for all x and y in S . As e is left identity in S . So, $f(xy) = f((ex)y) \geq f(x)$, which implies that $f(xy) \geq f(x) \wedge f(y)$ for all x and y in S . Thus f is fuzzy AG-subgroupoid. Consider, for any x, y and z in S , $f((xy)z) = f((x(ey))z) = f((e(xy))z) \geq f(xy) = f((ex)y) \geq f(x)$. Also $f((xy)z) = f((zy)x) = f((z(ey))x) = f((e(zy))x) \geq f(zy) = f((ez)y) \geq f(z)$. Hence $f((xy)z) \geq f(x) \wedge f(z)$ for all x, y and z in S . $\square$

Proposition 6. *Let f is a fuzzy subset of an AG-groupoid S with left identity. If f is a fuzzy left(right, two-sided) ideal in S then $f \circ f$ is a fuzzy ideal in S .*

Proof. Let f is a fuzzy left ideal in an AG-groupoid S , then by lemma 1 $S \circ f \subseteq f$. By use of lemma 3 and corollary 1, $S \circ (f \circ f) = (S \circ S) \circ (f \circ f) = (S \circ f) \circ (S \circ f) \subseteq f \circ f$. Also by proposition 1, $(f \circ f) \circ S = (S \circ f) \circ f \subseteq f \circ f$. If f is a fuzzy right ideal in S then by lemma 5 f is a fuzzy left ideal. $\square$

Corollary 3. *Let f is a fuzzy subset of an AG-groupoid S with left identity. If f is a fuzzy left ideal in S then $f \circ f$ is a fuzzy bi-ideal and an interior ideal in S .*

Proof. By proposition 6, $f \circ f$ is a fuzzy ideal in S . Now by lemmas 11 and 12, $f \circ f$ is a fuzzy interior and fuzzy bi-ideal of S . $\square$

Theorem 6. *In an AG-groupoid S , every fuzzy ideal is a fuzzy bi-ideal and an interior ideal of S .*

Proof. Let f be a fuzzy ideal of an AG-groupoid S . Clearly f is AG-subgroupoid of S by lemma 1, since $f \circ f \subseteq S \circ f \subseteq f$. Consider, $(f \circ S) \circ f \subseteq f \circ f \subseteq f$, which by lemma 7 shows that f is a bi-ideal in S . Now, consider $(S \circ f) \circ S \subseteq f \circ S \subseteq S$, which by lemma 10 shows that f is an interior ideal in S . $\square$

A fuzzy ideal f of an AG-groupoid S is called fuzzy prime ideal if for any two fuzzy ideals g and h of S , $g \circ h \subseteq f$ implies that either $g \subseteq f$ or $h \subseteq f$. An AG-groupoid S is fully fuzzy prime if every fuzzy ideal is prime in S . A fuzzy left ideal f of an AG-groupoid S is called fuzzy quasi-prime if for any two fuzzy left ideals g and h of S , $g \circ h \subseteq f$ implies that either $g \subseteq f$ or $h \subseteq f$. An AG-groupoid S is fully fuzzy quasi-prime if every fuzzy left ideal is quasi-prime in S . A fuzzy left ideal f of an AG-groupoid S is called fuzzy semiprime left ideal of S if for any fuzzy left ideal g of S , $g^2 \subseteq f$ implies $g \subseteq f$.

The collection of fuzzy subsets $F(S)$ of an AG-groupoid S is totally ordered if for all fuzzy ideals f, g of S either $f \subseteq g$, or $g \subseteq f$.

Theorem 7. *An AG-groupoid S with left identity is fully fuzzy prime if and only if every fuzzy ideal is idempotent and fuzzy ideals are totally ordered.*

Proof. Let S is fully fuzzy prime. Let f is fuzzy ideal of S , then since $f \circ f \subseteq S \circ f \subseteq f$. By proposition 6 $f \circ f$ is fuzzy ideal and by hypothesis $f \circ f$ is prime. So, $f \circ f \subseteq f \circ f$, implies that $f \circ f \subseteq f$. Consider, f and g be fuzzy ideals of S , then $f \circ g \subseteq f \circ S \subseteq f$ and also $f \circ g \subseteq S \circ g \subseteq g$. Hence, $f \circ g \subseteq f \cap g$, where $f \cap g$ is fuzzy ideal by lemma 2. By definition of fully fuzzy prime $f \subseteq f \cap g$ or $g \subseteq f \cap g$, which implies that $f \subseteq g$ or $g \subseteq f$. Conversely, let every fuzzy ideal of S is idempotent and fuzzy ideals are totally ordered. Let f be any fuzzy ideal of S such that $g \circ h \subseteq f$ where f and g are fuzzy ideals of S . Since ideals are totally ordered so for g and h either $g \subseteq h$ or $h \subseteq g$. Let $g \subseteq h$. Since g is idempotent so, $g = g \circ g \subseteq g \circ h \subseteq f$. Similarly, for $h \subseteq g$ we have, $f \subseteq g$. $\square$

Proposition 7. *Let S be an AG-groupoid with left identity. If S is fully fuzzy quasi-prime then every fuzzy left ideal is idempotent.*

Proof. Let f be a fuzzy left ideal in an AG-groupoid with left identity S , where S is fully fuzzy quasi-prime. Now, $S \circ f \subseteq f$ but $f \circ f \subseteq S \circ f$ so $f \circ f \subseteq f$. By proposition 6 $f \circ f$ is fuzzy ideal and by hypothesis $f \circ f$ is quasi-prime. So, $f \circ f \subseteq f \circ f$, implies that $f \circ f \subseteq f$. $\square$

Theorem 8. *Let S be an AG-groupoid with left identity. If S is fully fuzzy quasi-prime then for every fuzzy left ideals f and g of S , $f \circ g = f \cap g$.*

Proof. Let f and g are fuzzy left ideals in an AG-groupoid with left identity S , where S is fully fuzzy quasi-prime. So, $S \circ f \subseteq f$ and $S \circ g \subseteq g$, which implies that $g \circ f \subseteq f$ and $f \circ g \subseteq g$. By proposition 7 and theorem 1, $g \circ f = (g \circ g) \circ (f \circ f) = (f \circ f) \circ (g \circ g) = f \circ g$, which implies that $f \circ g \subseteq f \cap g$. Since $f \cap g \subseteq f$ and $f \cap g \subseteq g$ so, $(f \cap g) \circ (f \cap g) \subseteq f \circ g$. By lemma 2 and proposition 7, $f \cap g$ is fuzzy left ideal and hence idempotent. So, $f \cap g \subseteq f \circ g$ which implies that $f \cap g = f \circ g$. $\square$

Corollary 4. *The set of fuzzy quasi-prime ideals of AG-groupoid with left identity forms a semilattice structure.*

Proof. It is an easy consequence. $\square$

Theorem 9. *Let S be an AG-groupoid with left identity, then followings are equivalent:*

- (i) each left ideal of S is idempotent,
- (ii) each fuzzy left ideal of S is idempotent,
- (iii) for each pair of fuzzy left ideals f, g of S , $f \circ g = f \cap g$,
- (iv) each fuzzy left ideal of S is fuzzy semiprime left ideal.

Proof. (i) $\Leftrightarrow$ (ii) from [9].

Let (ii) holds that is every fuzzy left ideal of S is idempotent. Let f and g are fuzzy left ideals which are idempotent, then $f \circ g \subseteq g$ and $g \circ f \subseteq f$. But $f \circ g = (f \circ f) \circ g = (g \circ f) \circ f \subseteq f \circ f \subseteq f$ and so $f \circ g \subseteq f \cap g$. Also $f \cap g \subseteq f$ and $f \cap g \subseteq g$ which implies that $(f \cap g) \circ (f \cap g) \subseteq f \circ g$. We know that intersection of fuzzy left ideals f and g is fuzzy left ideal so $f \cap g$ is fuzzy left ideal of S and hence an idempotent, which implies that $f \cap g \subseteq f \circ g$.

(iii) $\Rightarrow$ (iv) is same as in [9] using the hypothesis for fuzzy left ideal f in S that $f \circ f = f \cap f = f$.

Let (iv) holds that each fuzzy left ideal f is semiprime in S . By use of theorem 1 and lemma 4, $S \circ (f \circ f) = f \circ (S \circ f) = f \circ f$ that is $f \circ f$ is fuzzy left ideal of

S and hence semiprime. Since $f \circ f \subseteq f \circ f$ which implies that $f \subseteq f \circ f$. But we know that f is fuzzy left ideal so $f \circ f \subseteq S \circ f \subseteq f$. $\square$

REFERENCES

- [1] Mordeson, J. N., *Fuzzy semigroups*, Springer-Verlag Berlin Heidelberg, 2003.
- [2] Kuroki, N., On fuzzy ideals and fuzzy bi-ideals in semigroups, *Fuzzy Sets and Systems*, 5 (1981) 203-215.
- [3] Mushtaq, Q. and Madad Khan, Ideals in AG-band and AG*-groupoid, *Quasigroups and Related Systems*, 14 (2006), 207-215.
- [4] Xiang-Yun Xie, On prime, quasi-prime, weakly quasi-prime fuzzy left ideals of semigroups, *Fuzzy Sets and Systems*, 123 (2001), 239-249.
- [5] Xiang-Yun Xie and Jian Tang, Fuzzy radicals and prime fuzzy ideals of ordered semigroups, *Information Sciences*, 178 (2008), 4357-4374.
- [6] Kazim, M. A. and M. Naseeruddin, On LA-Semigroups, *Allig. Bull. Math.*, 8 (1972), 1-7.
- [7] Protic, P. V. and N. Stevanović, On Abel-Grassmann's groupoids, *Proc. Conf. Pristina*, 1994, 31-38.
- [8] Wang Xue-ping, Mo Zhi-wen and Liu Wang-jin, Fuzzy Ideals Generated by Fuzzy point in Semigroup, *J. Sichuan Normal Univ.*, 15 (1992) 17-24.
- [9] Shabir, M., Fully fuzzy prime semigroups, *Inter. J. of Math. and Mathematical Sciences*, 2005 (2005), 163-168.
- [10] Zadeh, L.A., Fuzzy sets, *Inform. Control*, 8 (1965), 338-353.
- [11] Rosenfeld, A., Fuzzy groups, *J. Math. Anal. Appl.*, 35 (1971), 512-517.
- [12] Kuroki, N., Fuzzy bi-ideals in Semigroups, *Comment. Math. Univ. St. Pauli*, 27 (1979), 17-21.

COMSATS INSTITUTE OF INFORMATION TECHNOLOGY, ABBOTTABAD.

E-mail address: madadmath@yahoo.com, mailnouman@gmail.com